

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/10, C07K 1/14</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 97/05248</b> <b>(43) International Publication Date:</b> 13 February 1997 (13.02.97)
<b>(21) International Application Number:</b> PCT/US96/11875 <b>(22) International Filing Date:</b> 18 July 1996 (18.07.96) <b>(30) Priority Data:</b> 08/509,164                      31 July 1995 (31.07.95)                      US <b>(71)(72) Applicant and Inventor:</b> CHOMCZYNSKI, Piotr [US/US]; 778 Avon Fields Lane, Cincinnati, OH 45229 (US). <b>(74) Agents:</b> AHRENS, Gregory, F. et al.; Wood, Herron & Evans, P.L.L., 2700 Carew Tower, Cincinnati, OH 45202 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> SOLUTION CONTAINING CHAOTROPIC AGENT AND PROCESS USING IT FOR ISOLATION OF DNA, RNA AND PROTEINS  <b>(57) Abstract</b>  Solutions and methods are disclosed for the effective, simple isolation/extraction of DNA, RNA and proteins from a single biological material sample, such as cells, tissues and biological fluids. The preferred solutions include effective amounts of a chaotropic agent(s), buffer, reducing agent, and may or may not include an organic solvent. Genomic DNA and total RNA can be isolated utilizing the solutions and methods of the invention in as little as 20 minutes, and proteins in as little as 30 minutes.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

-1-

**SOLUTION CONTAINING CHAOTROPIC AGENT AND PROCESS USING IT FOR ISOLATION OF  
DNA, RNA AND PROTEINS****Field of the Invention**

The invention relates to compositions and methods for isolating nucleic acids (both RNA and DNA) and proteins from biological materials. And more particularly, the invention relates to nucleic acid and protein isolation methods employing non-toxic chaotropic agents.

**Background of the Invention**

The continuous advances in molecular biology, biotechnology and clinical research have resulted in an ever increasing number of uses for DNA, RNA and proteins. For example, polymerase chain reaction (PCR) technology has dramatically expanded the use of DNA and RNA in basic research, in clinical diagnostics such as detection of messenger RNA (mRNA) by reverse transcription PCR (RT-PCR), and the use of PCR in detection of genetic defects. In the protein field, identification of proteins by Western blotting has become an important tool in studying gene expression in basic research and identification of specific proteins for diagnostic purposes, as exemplified by viral protein detection.

-2-

The increased use of RNA, DNA and proteins has created a need for fast, simple and reliable methods and reagents for isolating DNA, RNA and proteins. In many applications, collecting the biological material sample and subsequent analysis thereof would be substantially simplified if the three cellular components (RNA, DNA and proteins) could be simultaneously isolated from a single sample. The simultaneous isolation is especially important when the sample size is so small, such as in biopsy, that it precludes its separation into smaller samples to perform separate isolation protocols for DNA, RNA and proteins.

There are known methods for isolating DNA, RNA and proteins from a single biological material sample. One such method is described in Coombs, L. M., et al.: "Simultaneous Isolation of DNA, RNA and Antigenic Protein Exhibiting Kinase Activity from Small Tumor Samples Using Guanidine Isothiocyanate", Anal. Biochem., 188, 338-343 (1990). The Coombs et al. method is based on ultracentrifugation of the sample homogenate in a guanidine-cesium chloride solution. The sample is homogenized in 4 M guanidine thiocyanate and then overlaid on a cesium chloride (CsCl) solution and centrifuged at  $>100,000g$  for about 18 hours. Following centrifugation, DNA, RNA and proteins are separated and purified over the next 12-24 hours. This method has several limitations or drawbacks, including the prolonged time required for isolation and the limited number and size of samples which can be processed with an ultracentrifuge. Also, the high cost of an ultracentrifuge may be prohibitive in certain circumstances.

-3-

Another method for the simultaneous isolation of DNA, RNA and proteins from a single biological material sample is the subject of my earlier U.S. Patent No. 5,346,994 (the '994 patent). That method is based on liquid-phase separation using phenol and guanidine thiocyanate. A biological sample is homogenized in the aqueous solution of phenol and guanidine thiocyanate and the homogenate thereafter is mixed with chloroform. Following centrifugation, the homogenate separates into an organic phase, an interphase and an aqueous phase. Proteins are sequestered into the organic phase, DNA into the interphase and RNA into the aqueous phase. Next, each component is precipitated from the corresponding phase using ethanol and is then washed. The whole procedure can be completed in about 2-3 hours, and is especially useful for the isolation of high quality RNA from a variety of sources. One drawback to this method is the use of highly toxic phenol.

There are many known methods for the separate isolation of DNA, RNA and proteins from biological material; i.e., protocols for isolating a single one of these components from a sample. In typical DNA isolation methods, a biological sample is lysed in a lysing solution and then the DNA is isolated from the lysate according to any one of a variety of multi-step protocols, which may take from one hour to several days to complete. Frequently recommended DNA isolation methods involve the use of toxic phenol. See, Sambrook, J. et al., "Molecular Cloning", Vol. 2, pp. 9.14-9.23, Cold Spring Harbor Laboratory Press (1989) and Ausubel, Frederick M. et al., "Current Protocols in Molecular Biology", Vol. 1, pp. 2.2.1-2.4.5, John Wiley & Sons, Inc. (1994). Typically, a biological sample is lysed in a detergent

-4-

solution and the protein component of the lysate is digested with proteinase for 12 - 18 hours. Next, the lysate is extracted with phenol to remove most of the cellular components, and the remaining aqueous phase is processed further to isolate DNA. In another method, described in Van Ness et al. U.S. Patent 5,130,423, non-corrosive phenol derivatives are used for the isolation of nucleic acids. The resulting preparation is a mix of RNA and DNA.

DNA isolation methods utilizing non-corrosive chaotropic agents have also been developed. These methods, which are based on the use of guanidine salts, urea and sodium iodide, involve lysis of a biological sample in a chaotropic aqueous solution and subsequent precipitation of the crude DNA fraction with a lower alcohol. The final purification of the precipitated, crude DNA fraction can be achieved by any one of several methods, including column chromatography as described in "RapidPrep™ Genomic DNA Isolation Kits For Cells and Tissue: Versatility at Your Fingertips!", Analects, Vol 22, No. 4, Pharmacia Biotech, 1994, or exposure of the crude DNA to a polyanion-containing protein as described in Koller U.S. Patent 5,128,247.

Yet another method of DNA isolation, which is described in Botwell, D.D.L., "Rapid Isolation of Eukaryotic DNA", Anal. Biochem. 162, 463-465 (1987) involves lysing cells in 6 M guanidine hydrochloride, precipitating DNA from the lysate at acid pH by adding 2.5 volumes of ethanol, and washing the DNA with ethanol. It is believed that the resulting DNA may be contaminated, however, with a low molecular weight material such as RNA and pigments. This conclusion is in agreement with the well known report showing that under similar conditions RNA can be precipitated from cell or

-5-

tissue lysate. See Chirgwin, J.M. et al., "Isolation of Biologically Active Ribonucleic Acid from Sources Enriched in Ribonuclease", Dept. of Biochemistry and Biophysics, Univ. of California, Vol. 18, No. 24, pp. 5294-5299 (1979).

5

Sodium iodide, another chaotropic agent, has been used in DNA isolation, but its use requires an additional purification step consisting of adsorption of the isolated DNA on glass beads. Although this method is relatively simple, it results in a low yield of isolated DNA.

10

In still another approach, the bulk of cytoplasmic proteins and RNA are removed by lysing cell samples in a detergent solution. The lysate is then fractionated into the nuclear and cytoplasmic fractions. And thereafter, the DNA is purified by dissolving the nuclear fraction in a chaotropic solution, precipitating and washing with ethanol. This method, described in Ciulla, T.A. et al., "A Simple Method for DNA Purification from Peripheral Blood", Anal. Biochem. 174, 485-488 (1988), can be completed in about 2 hours and is useful for isolating DNA from whole blood.

15

20

Known techniques for isolating RNA typically utilize either guanidine salts or phenol extraction, as described for example in Sambrook, J. et al., "Molecular Cloning", Vol. 1, pp. 7.3-7.24, Cold Spring Harbor Laboratory Press (1989) and Ausubel, F. M. et al., "Current Protocols in Molecular Biology", Vol. 1, pp. 4.0.3-4.4.7, John Wiley & Sons, Inc. (1994). Phenol-based techniques are multi-step procedures requiring several hours or days to complete. Similarly, the guanidine-based RNA isolation methods require at least several hours and take many steps to complete. In my earlier

-6-

U.S. Patent No. 4,843,155, phenol and guanidine procedures were uniquely combined, resulting in a simple method of total RNA isolation which can be completed in 3 hours. The method of the '155 patent was further improved upon in my '994 patent, which allows for completion of the RNA isolation in about 1 hour.

There are also known techniques for the simultaneous isolation of DNA and RNA, as referenced in my earlier '994 patent, the disclosures of which are incorporated herein by reference. All of these techniques utilize phenol extraction as a necessary step for isolating RNA and DNA free of protein contamination.

Heretofore, it has been the commonly accepted view that precipitation of nucleic acids from chaotropic solvents does not result in pure nucleic acid preparations. Contrary to this view, however, the present inventor has found that under certain conditions, as described in full detail herein, the use of chaotropic agents alone will result in isolation of assay ready, high quality DNA and RNA. This unexpected finding led to the development of a very simple, effective method and product for the simultaneous isolation of DNA, RNA and proteins from a single sample for subsequent use in molecular biology, biotechnology, clinical research and other applications.

#### Summary of the Invention

The products and methods of the present invention provide a highly effective, simple means of extracting DNA, RNA and proteins from a single biological material sample, such as cells, tissues and biological fluids. Advantageously, these results can be achieved without the use of toxic or



-7-

corrosive reagents and without the use of expensive ultracentrifugation equipment. Genomic DNA and total RNA can be isolated utilizing the products and methods of the invention in as little as 20 minutes, and proteins in as little as 30 minutes. These results are substantially faster than existing methods for the simultaneous isolation of DNA, RNA and proteins. The invention is also applicable to the separate isolation of RNA only, DNA only, proteins only or any combination of two of these cellular components. The resulting genomic DNA and total RNA isolated utilizing the methods and products of the invention are of high quality suitable for use in research, biotechnology, etc. The invention is in part based on the unexpected finding that utilizing the products of the invention, RNA is precipitated prior to DNA, which is contrary to prior art methodologies. In particular, the finding that the solvent solution of the present invention precipitates RNA prior to DNA is in striking contrast to the described prior art such as Koller U.S. Patent No. 5,128,247, wherein it describes that DNA exhibits a lower solubility than RNA and apparently can be precipitated more easily than RNA. Furthermore, substantially lower amounts of organic solvents are required to effect the precipitation of the cellular components.

In its broadest aspects, the invention encompasses solutions for isolating substantially pure and undegraded RNA, DNA and protein from biological materials, including tissue, cells and fluids. The solution preferably includes effective amounts of a chaotropic agent(s), buffer, reducing agent, and water (with or without organic solvent(s)). The chaotropic agent(s) act to dissociate proteins from nucleic acids (RNA and DNA) and inhibit the activity

-8-

of nucleic acid degradation enzymes. The actions of the chaotropes are potentiated by the reducing agent present in the solvent solution. When the solution includes an organic solvent, its presence prevents solubilization of RNA, thereby making it possible to remove RNA from the homogenate formed when the tissue sample is homogenized in the solvent solution by a brief centrifugation step. When the solution does not include an organic solvent, the RNA is precipitated from the homogenate subsequent to homogenization by the addition of an organic solvent.

Preferred chaotropic agents for the solution include guanidine thiocyanate, guanidine hydrochloride, and mixtures thereof. These components may be supplemented with other chaotropes, such as urea or sodium iodide. The preferred concentration of chaotropes in the solution is in the range of about 2 M - 7 M. Preferably, the reducing agent is non-toxic, such as 2-aminoethanethiol. If required, this can be substituted with 2-mercaptoethanol; however, this is a toxic composition. The reducing agent facilitates denaturation of RNase by the chaotropes and aids in the isolation of undegraded RNA.

The solution of the present invention preferably contains a sufficient amount of buffer to maintain the pH of the solution above about 6. For the simultaneous isolation of RNA, DNA and proteins, the pH should be maintained in the range of about 6-7.5. For the isolation of DNA alone, the effective pH range may be about 6-12. The preferred buffers for use in the solutions of the invention include tris-hydrochloric acid, sodium phosphate, sodium acetate, sodium tetraborate-boric acid and glycine-sodium hydroxide.

-9-

As stated, the solution may contain an organic solvent. The preferred solvents are lower alcohols such as methanol, ethanol and isopropanol. However, other water miscible solvents can be used which will achieve the desired effect, such as acetone, polyethylene glycol and dimethylsulfoxide, or mixtures of any of the foregoing. The effective concentration of organic solvents in the product of the invention is in the range of about 15-30% by volume.

The solution of the invention may contain additional components, including organic and inorganic salts such as sodium chloride, potassium chloride, ammonium chloride, sodium phosphate, sodium acetate, sodium nitrite, lithium chloride, and sodium bromide. Furthermore, compatible detergents such as sarcosines and polyoxyethylenesorbitan, and chelating agents such as ethylenediamine tetraacetic acid and citric acid can be utilized to promote tissue solubilization and precipitation of nucleic acids.

In another aspect, the invention encompasses methods of isolating substantially pure RNA, DNA and proteins from biological material samples. Utilizing the solutions of the present invention, and particularly those wherein the solution includes an organic solvent, the method includes an initial step of homogenizing a biological material sample in the solution to form an homogenate. The presence of the organic solvent prevents solubilization of RNA, thereby making it possible to remove RNA from the homogenate by brief centrifugation (sedimentation). Thereafter, DNA is precipitated from the remaining homogenate by adding an effective amount of an organic solvent and recovering the precipitated DNA by brief centrifugation (sedimentation).

-10-

Finally, adding an effective amount of an organic solvent to the remaining homogenate precipitates proteins therefrom. The successive addition of organic solvents to precipitate genomic DNA and proteins from the post-RNA homogenate require a limited amount of the organic solvent. Following a wash with ethanol, the three cellular components are each separately and fully ready for use in molecular biology, biotechnology and clinical research applications. Preferably, the organic solvent utilized to precipitate DNA is added in a ratio of about 0.15-0.3 volumes of solvent per one volume of initial homogenate. The organic solvent added to precipitate proteins is preferably added in the ratio of about 3-4 volumes of solvent per one volume of initial homogenate. One preferred organic solvent for this purpose is isopropanol, although other suitable organic solvents can be used.

In an alternative methodology, wherein the solution does not include an organic solvent component, the homogenate is formed and an initial centrifugation step is performed to remove any unhomogenized tissue. Thereafter an effective amount of an organic solvent is added to preclude the solubilization of total RNA. This makes it possible to recover the RNA from the homogenate by a brief centrifugation (sedimentation). Preferably, the organic solvent added to precipitate RNA is added in a ratio of about 0.15-0.3 volumes of solvent per one volume of initial homogenate. Once separated, the RNA is washed with ethanol and the RNA precipitate may be dissolved in formamide and stored at -20°C. The use of formamide as a solubilization agent is beneficial in that it protects RNA from degradation by RNase, which may otherwise contaminate the isolated RNA.

-11-

Following precipitation of the RNA, DNA is precipitated by adding an additional 0.15-0.3 volumes of organic solvent per one volume of the initial homogenate. The DNA precipitate is removed from the homogenate by spooling or brief centrifugation (sedimentation). Proteins and other cellular components are retained in the homogenate and the protein component is precipitated by the addition of 3-4 volumes of an organic solvent per one volume of the initial homogenate.

Overall, and in comparison with other known methodologies, the precipitation of nucleic acids according to the present invention is performed with a substantially reduced amount of organic solvents. For example, in Sambrook, J. et al., "Molecular Cloning", Vol. 1, pp. 7.3-7.24, Cold Spring Harbor Press (1989), the precipitation of RNA requires the addition of 0.5-2.5 volumes of a lower alcohol added to one volume of the chaotropic solution. In addition, the RNA precipitation was carried out for several hours at -20-4°C. In contrast, the precipitation of RNA according to the present method is completed in about 3-5 minutes and is performed at room temperature. With respect to the DNA precipitation, Botwell, D.D.L. "Rapid Isolation of Eukaryotic DNA", Anal. Biochem. 162, 463-465 (1987), describes precipitation from chaotropic solutions carried out with 2-2.5 volumes of ethanol. Likewise, high volumes of alcohol have been recommended for the effective DNA precipitation from non-chaotropic solutions, as exemplified in Ausubel, F.M. et al., "Current Protocols in Molecular Biology", Vol. 1, pp. 221-245, John Wiley & Sons, Inc. (1994).

-12-

The unexpectedly low concentration of organic solvents required for precipitating RNA and DNA from the chaotropic solutions of the present invention makes it feasible to obtain RNA and DNA in highly pure form for use in molecular biology and biotechnology, as well as clinical research and other applications.

These and other features and advantages of the present invention will become apparent to persons skilled in the art upon review of the detailed description and working examples herein.

#### Brief Description of the Drawings

Fig. 1A shows the electrophoretic analysis of nucleic acids isolated in accordance with Example 1 herein;

Fig. 1B shows the utility of the total RNA in RT-PCR;

Fig. 1C shows the utility of the genomic DNA in PCR;

Fig. 1D shows the results of protein analysis by Western blotting;

Fig. 2A shows the electrophoretic analysis of nucleic acids isolated in accordance with Example 2 herein;

Fig. 2B shows the results of total RNA analysis by Northern blotting;

Fig. 2C shows the results of genomic DNA analysis by Southern blotting;

Fig. 2D shows the results of protein analysis by Western blotting; and

Fig. 3 shows the results of genomic DNA analysis by electrophoresis as isolated in Example 5.

Detailed Description of the Invention

Preferred solutions and methods according to the present invention are described in the following working Examples.

Example 1 - Simultaneous Isolation of DNA, RNA & Proteins From Cells

5 A method of the invention was used to simultaneously isolate DNA, RNA and proteins from rat somatomammotroph P0 cells. About  $10^8$  P0 cells were lysed (homogenized) in a 10ml of solution of the invention, which contained: 4 M guanidine thiocyanate (Amresco, Inc., Solon, OH), 17% isopropanol, 0.1 M sodium acetate, 0.1 M 2-aminoethanethiol hydrochloride  
10 (Sigma, St. Louis, MO) and 0.2% sarkosyl in water. The solution was adjusted to pH 7.0 with hydrochloric acid. Unless stated otherwise, chemical reagents were obtained from Fisher Scientific (Pittsburgh, PA). Next, 0.6 ml aliquots of the lysate (homogenate) were frozen or used immediately for the isolation. The 0.6 ml aliquot utilized as described below contained  $37.3 \mu\text{g}$  DNA, as  
15 determined by the diphenylamine reaction.

RNA isolation. The lysate (0.6 ml) was centrifuged for 8 minutes at 10,000 g at room temperature to sediment total RNA. The post-RNA lysate was transferred to a fresh tube and saved for the DNA and protein isolation described below. The RNA pellet was washed with 1 ml of 95% ethanol by  
20 vortexing and pipetting off the ethanol. Finally, RNA was dissolved in formamide and stored at  $-20^\circ\text{C}$ . The total RNA isolation was completed in 11 minutes. The isolated RNA exhibited a 260/280 ratio of  $1.79 \pm 0.05$ , with a yield of  $49.2 \pm 2.8 \mu\text{g}$  RNA (mean  $\pm$  SD,  $n=3$ ). Northern blotting of the

isolated RNA preparations showed an undegraded pattern of mRNA when tested for growth hormone, prolactin,  $\beta$ -actin and GAPDH mRNAs.

For use in the reverse transcription PCR (RT-PCR), an aliquot of the RNA-solubilized formamide was precipitated with 3 volumes of ethanol.

5 The precipitate was dissolved in water and used for RT-PCR.

It has been found that spectrophotometrical measurements to determine the optical density of RNA and DNA are substantially improved when performed in a solution containing higher than usual concentrations of chelating agents. For this purpose, the concentration of chelating agent(s) should be  
10 higher than 5 mM, with the optimum for ethylenediamine tetraacetate (EDTA) at 10 mM and for citrate at 30 mM. This is a new and unexpected finding. Typically, optical density readings of RNA and DNA are performed in water or 1 mM EDTA.

At higher concentration of chelating agents, the optical density  
15 readings are more reproducible and result in a higher 260/280 ratio. For example, the RNA preparation described in Example 1 had a 260/280 ratio of  $1.79 \pm 0.05$  when measured in water, and a ratio of  $1.97 \pm 0.01$  when measured in 10 mM EDTA.

DNA isolation. DNA was precipitated from the post-RNA lysate  
20 by the addition of 0.15 ml of isopropanol. The floating DNA precipitate was swirled (spooled) onto the pipette tip and transferred to a new tube. The remaining post-DNA lysate was saved for the protein isolation described below. The DNA was washed by mixing it with 1 ml of 95% ethanol and pipetting out the ethanol wash. The final DNA preparation was dissolved in 8 mM NaOH by



-15-

gentle pipetting followed by neutralization of the solution with N-[2-hydroxyethyl] piperazine-N-[2-ethane sulfinic acid] (HEPES, free acid). The isolation was completed in less than 14 minutes. The isolated DNA exhibited a 260/280 ratio  $1.81 \pm 0.02$  (SD, n=3), indicating a lack protein contamination.

5 The average yield from three isolations was  $33.9 \pm 2.9$  (SD, n=3)  $\mu$ g DNA. As compared with the original amount of DNA in the lysate, determined by the diphenylamine reaction, the method of the invention provided 91 % recovery of DNA.

Protein isolation. Proteins were precipitated from the post-DNA  
10 lysate by the addition of 1.8 ml of isopropanol and centrifugation at 10,000 g for 5 minutes. The precipitate was washed with 95 % ethanol and dissolved in 0.1 N acetic acid. Alternatively, the precipitate could be dissolved in 0.1 % sodium dodecylsulphate, or in water. The protein isolation was completed in 22 minutes. The isolated protein preparation was tested by Western blotting using  
15 a specific anti-rat prolactin antibody. The presence of the prolactin specific band in Western blotting is indicative of the good quality of the isolated protein preparation.

The results of tests performed with the simultaneously isolated RNA, DNA and proteins are shown in Figure 1. Figure 1 A shows the results  
20 of nucleic acids electrophoresed in 1 % agarose gel and stained with ethidium bromide. Lanes 1 and 2 show undegraded total RNA (3  $\mu$ g/lane); lanes 3 and 4 show high molecular weight genomic DNA (3  $\mu$ g/lane); and lanes 5 and 6 show genomic DNA (3  $\mu$ g) digested for 2 hours with EcoR1 restrictase. As the results demonstrate, there is no detectable DNA in the RNA preparation and no

-16-

detectable RNA in the DNA preparation. The total digestion of DNA by EcoR1  
restrictase is indicative of the good quality of the isolated DNA. Figures 1B-D  
show the results of RT-PCR, PCR and Western blotting performed with total  
RNA, genomic DNA and proteins, respectively, isolated as described  
hereinabove. The RT-PCR (Fig. 1B) and PCR (Fig. 1C) were performed using  
glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and rat growth hormone  
(GH) primers, respectively, and Western blotting (Fig. 1C) was performed  
using anti-rat prolactin antibody. Amplification of the 374 base pair GAPDH  
DNA fragment in RT-PCR and the 686 base pair GH DNA fragment in PCR  
indicates that the isolated nucleic acid preparations (RNA and DNA) are  
adequately purified for the PCR reaction. Also, the presence of a prolactin  
specific band in Western blotting shows the high quality of the isolated protein  
preparation.

Similar results in the simultaneous isolation of RNA, DNA and  
proteins have been obtained when, in place of isopropanol, other water miscible  
organic solvents such as ethanol, methanol, acetone, dimethylsulfoxide,  
polyethylene glycol or mixtures of these solvents are used. These water miscible  
organic solvents can be used as components of the lysing solution and/or as the  
precipitating agents. All solvents are available from companies such Aldrich  
Chemical Co., Inc. (Milwaukee, WI) or Fluka Chemical Corp. (Ronkonkoma,  
NY).

#### EXAMPLE 2 - Simultaneous Isolation of RNA, DNA and Proteins From Tissues.

In this embodiment of the method, the solution does not include  
an organic solvent component. This permits the removal of any unhomogenized

-17-

tissue fragments from the lysate (homogenate) by a brief initial centrifugation. Thereafter, RNA is precipitated from the clear lysate by the addition of 0.2-0.3 volumes of an organic solvent.

5 A frozen sample of rat pituitary was homogenized in a hand held glass-Teflon homogenizer with 1 ml of a lysing solution containing 4 M guanidine thiocyanate, 0.1 M sodium acetate, 0.1 M 2-aminoethanethiol hydrochloride and 0.2% sarkosyl in water. The solution was adjusted to pH 7.0 with hydrochloric acid. The homogenate was centrifuged at 10,000 g for 5 minutes. The clear supernatant was transferred to a new tube, mixed with 0.3  
10 ml of isopropanol and stored for 3 minutes at room temperature to precipitate RNA. The precipitated RNA was removed by centrifugation at 10,000 g for 8 minutes and dissolved in formamide. The post-RNA supernatant was further processed to isolate DNA and proteins in the same manner as described in Example 1. DNA and proteins were successively precipitated from the post-  
15 RNA lysate by the addition of 0.5 ml acetone and 10.75 ml of acetone, respectively. The isolated RNA exhibited a 260/280 ratio of 1.74 and the yield was 0.06 mg. The DNA exhibited a 260/280 ratio of 1.78 and the yield was 0.04 mg. As shown in Figure 2A, there was no detectable DNA present in the RNA preparation and no detectable RNA present in the DNA preparation. As  
20 in Fig. 1A, the nucleic acids were electrophoresed in 1% agarose gel and stained with ethidium bromide. Lanes 1 and 2, total RNA (3  $\mu$ g/lane); lanes 3 and 4, genomic DNA (3  $\mu$ g/lane); and lanes 5 and 6, genomic DNA (3  $\mu$ g) digested for 2 hours with EcoR1 restrictase.

-18-

The preparations of total RNA, genomic DNA and proteins were tested by Northern, Southern and Western blotting for the rat growth hormone (GH) mRNA, GH gene, and GH, respectively. All three analyses, depicted in Figs. 2B-D, respectively, show that the method of invention yielded high quality RNA, DNA and protein preparations.

**EXAMPLE 3. Isolation of RNA From Cells without Centrifugation of The Initial Lysate**

Breast tumor MCF7 cells were grown in monolayer culture in a 3.5 cm petri dish. At the end of the culture period, the culture medium was removed and the cells were lysed by adding 1 ml of the lysing solution directly to the culture dish. The lysing solution used was that in Example 1. The lysate was centrifuged at 10,000 g for 5 minutes, and the RNA pellet was washed with 95% ethanol and dissolved in formamide. The isolated RNA exhibited a 260/280 ratio 1.81 and the yield was 22  $\mu$ g RNA.

**EXAMPLE 4. Isolation of RNA from Tissue with Centrifugation of The Initial Lysate to Remove Tissue Fragments**

Rat kidney (0.95 g) was homogenized in 19 ml of a lysing solution having the following composition: 4 M guanidine thiocyanate, 0.1 M sodium acetate and 0.2% sarkosyl in water. The solution was adjusted to pH 7.0 with hydrochloric acid. The homogenate was centrifuged at 10,000 g for 5 minutes to remove unsolubilized material. The resulting supernatant was transferred to a new tube and RNA was precipitated from the supernatant by the addition of 3.8 ml (0.3 volume) of isopropanol and centrifuged at 10,000 g for 8 minutes. The RNA pellet was dissolved in formamide and stored at -20 C.

-19-

The RNA preparation exhibited a 260/280 ratio 1.77 and the yield was 3.9 mg of RNA:

EXAMPLE 5 - Isolation of DNA from Cells without Centrifugation

5 The lysing solution used for this DNA isolation contained the following: 4 M guanidine thiocyanate, 0.1 M sodium acetate, 17% isopropanol, 0.2% sarkosyl in water. The lysing solution was adjusted to pH 9 by the addition of 0.4 M NaOH.

10 Rat pituitary PO cells were lysed in the lysing solution by repetitive pipetting. The lysate was mixed with 0.4 volume of ethanol and the precipitated DNA was spooled onto a pipette tip and washed twice with 95% ethanol. The resulting DNA was dissolved in 8 mM NaOH and neutralized to pH 8.0 with 0.1 M HEPES buffer.

15 This embodiment of the invention further shortens the DNA isolation protocol by omitting centrifugation of the lysate. This deletion of centrifugation results in only a minor contamination of the isolated DNA with RNA. Analysis of the genomic DNA isolated as described above was performed by electrophoreses thereof in 1% agarose gel and staining with ethidium bromide. Lanes 1 and 2 of Fig. 3 are genomic DNA (3  $\mu$ g/lane), and lanes 3 and 4 are genomic DNA digested for 2 hours with EcoR1. As is apparent in  
20 Figure 3, only a residual amount of RNA was detected in the low molecular weight region of the agarose gel by the ethidium bromide staining. Apparently, spooling DNA onto a pipette tip removes mainly DNA, while most of the partially hydrolyzed RNA (as indicated by its low molecular weight) remains in the lysate. The presence of a reducing agent is not necessary for the isolation

-20-

of DNA and the lysing solution can have pH within the range 6 - 12. The alkaline pH can be adjusted with NaOH, KOH or other organic or inorganic alkaline reagents. The best results are believed to be obtained when the lysate has a pH between 8 - 9.

5                   The unexpected and rapid hydrolysis of RNA occurring at pH 8 - pH 9 can be attributed to the presence of the chaotropic agent. This new finding allows for isolation of good quality DNA in a simple, one-step method.

10                   The protocol for DNA isolation without centrifugation can be completed in less than 10 minutes. This is believed to be the fastest and simplest method of genomic DNA isolation. Importantly, the only equipment required for this method are tubes and pipettes. This allows for performing the DNA isolation on a field trip or elsewhere with a limited access to laboratory equipment.

15                   **EXAMPLE 6 - Isolation of DNA from Tissues with One Centrifugation of the Lysate**

20                   Rat spleen (127 mg) was homogenized in a hand held glass-Teflon homogenizer with 5 ml of the lysing solution described in Example 5. The homogenate was centrifuged at 10,000 g for 5 minutes. The supernatant was transferred to a new tube and DNA was precipitated and washed as in Example 5. The isolated DNA exhibited a 260/280 ratio 1.76, and the yield was 1.93 mg DNA.

                  The scope of the present invention is not intended to be limited to the specific Examples described herein, but is to be accorded a scope commensurate with the appended claims.

-21-

What is claimed is:

-22-

1. A solution for isolating substantially pure and undegraded RNA, DNA and proteins from biological material, said solution comprising:  
effective amounts of  
at least one chaotropic agent, and  
a buffer present in an amount sufficient to maintain the pH  
of said solution in the range of about 6-12.
2. The solution of claim 1 wherein said chaotropic agent is selected from the group consisting of guanidine thiocyanate, guanidine hydrochloride, urea, sodium iodide and mixtures thereof.
3. The solution of claim 2 wherein said chaotropic agent is present at a concentration in the range of about 2M-7M.
4. The solution of claim 1 wherein said buffer is selected from the group consisting of tris-hydrochloric acid, sodium phosphate, sodium acetate, sodium citrate, sodium tetraborate and glycine-sodium hydroxide.
5. The solution of claim 1 wherein said buffer is present in an amount sufficient to maintain the pH of said solution in the range of about 6-7.5.
6. The solution of claim 1 further comprising at least one reducing agent.



-23-

7. The solution of claim 6 wherein said reducing agent is selected from the group consisting of 2-aminoethanethiol, 2-mercaptoethanol and mixtures thereof.
8. The solution of claim 1 further comprising a salt.
9. The solution of claim 8 wherein said salt is selected from the group consisting of sodium chloride, potassium chloride, ammonium chloride, sodium acetate, sodium nitrate, lithium chloride, sodium bromide and mixtures thereof.
10. The solution of claim 1 further comprising at least one chelating agent.
11. The solution of claim 10 wherein said chelating agent is selected from the group consisting of ethylenediamine tetraacetic acid and citric acid.
12. The solution of claim 1 further comprising a detergent.
13. The solution of claim 12 wherein said detergent is selected from the group consisting of sarcosine and polyoxyethylenesorbitan.

-24-

14. A solution for isolating substantially pure and undegraded RNA, DNA and proteins from biological material, said solution comprising:  
effective amounts of  
at least one chaotropic agent,  
a buffer present in an amount sufficient to maintain the pH  
of said solution in the range of 6-12, and  
an organic solvent present in an amount comprising about  
15-30% of said solution.
15. The solution of claim 14 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol, dimethylsulfoxide, and mixtures thereof.
16. The solution of claim 14 wherein said chaotropic agent is selected from the group consisting of guanidine thiocyanate, guanidine hydrochloride, urea, sodium iodide, and mixtures thereof.
17. The solution of claim 16 wherein said chaotropic agent is present at a concentration in the range of about 2M-7M.
18. The solution of claim 14 wherein said buffer is selected from the group consisting of tris-hydrochloric acid, sodium phosphate, sodium acetate, sodium citrate, sodium tetraborate and glycine-sodium hydroxide.

-25-

19. The solution of claim 14 wherein said buffer is present in an amount sufficient to maintain the pH of said solution in the range of about 6-7.5.
20. The solution of claim 14 further comprising at least one reducing agent.
21. The solution of claim 20 wherein said reducing agent is selected from the group consisting of 2-aminoethanethiol, 2-mercaptoethanol and mixtures thereof.
22. The solution of claim 14 further comprising a salt.
23. The solution of claim 22 wherein said salt is selected from the group consisting of sodium chloride, potassium chloride, ammonium chloride, sodium acetate, sodium nitrate, lithium chloride, sodium bromide and mixtures thereof.
24. The solution of claim 14 further comprising at least one chelating agent.
25. The solution of claim 24 wherein said chelating agent is selected from the group consisting of ethylenediamine tetraacetic acid and citric acid.

-26-

26. The solution of claim 14 further comprising a detergent.
27. The solution of claim 26 wherein said detergent is selected from the group consisting of sarcosine and polyoxyethylenesorbitan.
28. A method of isolating substantially pure and undegraded RNA, DNA and proteins from biological material, comprising the steps of:
- a) homogenizing a biological material sample in a solution of claim 1 to form an homogenate;
  - b) removing unhomogenized material from the homogenate by sedimentation;
  - c) precipitating RNA in the remaining homogenate by adding an effective amount of an organic solvent thereto, and recovering the precipitated RNA by sedimentation;
  - d) thereafter precipitating DNA in the remaining homogenate by adding an effective amount of organic solvent thereto, and recovering the precipitated DNA by one of spooling and sedimentation; and
  - e) thereafter precipitating proteins from the remaining homogenate by adding an effective amount of an organic solvent thereto, and recovering the precipitated proteins by sedimentation.
29. The method of claim 28 wherein said organic solvent added to precipitate RNA is added in a ratio of about 0.15-0.3 volumes of solvent per one volume of initial homogenate.

-27-

30. The method of claim 29 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol and dimethylsulfoxide.

31. The method of claim 28 wherein said organic solvent added to precipitate DNA is added in a ratio of about 0.15-0.3 volumes of solvent per one volume of initial homogenate.

32. The method of claim 31 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol and dimethylsulfoxide.

33. The method of claim 28 wherein said organic solvent added to precipitate proteins is added in a ratio of about 3-4 volumes of solvent per one volume of initial homogenate.

34. The method of claim 33 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol and dimethylsulfoxide.

-28-

35. A method of isolating substantially pure and undegraded RNA, DNA and proteins from biological material, comprising the steps of:

- a) homogenizing a biological material sample in the solution of claim 14 to form an homogenate;
- b) recovering substantially pure, undegraded RNA from the homogenate by sedimentation;
- c) thereafter precipitating DNA in the remaining homogenate by adding an effective amount of an organic solvent thereto, and recovering the precipitated DNA by one of sedimentation and spooling; and
- d) thereafter precipitating proteins from the remaining homogenate by adding an effective amount of organic solvent thereto, and recovering the precipitated proteins by sedimentation.

36. The method of claim 35 wherein said organic solvent added to precipitate DNA is added in a ratio of 0.15-0.3 volumes of solvent per one volume of the initial homogenate.

37. The method of claim 36 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol and dimethylsulfoxide.

38. The method of claim 35 wherein said organic solvent added to precipitate proteins is added in a ratio of 3-4 volumes of solvent per one volume of the initial homogenate.

-29-

39. The method of claim 38 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol and dimethylsulfoxide.

40. A method of isolating substantially pure and undegraded RNA from biological material, comprising the steps of:

a) homogenizing a biological material sample in a solution of claim 1 to form an homogenate;

b) removing unhomogenized material from the homogenate by sedimentation; and

c) precipitating RNA in the homogenate by adding an effective amount of an organic solvent thereto, and recovering the precipitated RNA by sedimentation.

41. The method of claim 40 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol and dimethylsulfoxide.

42. A method of isolating substantially pure and undegraded RNA from biological material, comprising the steps of:

a) homogenizing a biological material sample in a solution of claim 14 to form an homogenate; and

b) recovering substantially pure RNA from the homogenate by sedimentation.

-30-

43. A method of isolating substantially pure and undegraded DNA from biological material, comprising the steps of:

- a) homogenizing a biological material sample in a solution of claim 14 to form an homogenate; and
- b) precipitating DNA in the homogenate by adding an effective amount of an organic solvent thereto, and recovering the precipitated DNA by one of spooling and sedimentation.

44. The method of claim 43 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol and dimethylsulfoxide.

45. A method of isolating substantially pure and undegraded DNA from biological material, comprising the steps of:

- a) homogenizing a biological material in the solution of claim 1 to form an homogenate;
- b) removing unhomogenized material from the homogenate by sedimentation; and
- c) precipitating DNA in the homogenate by adding an effective amount of an organic solvent thereto, and recovering the precipitated DNA by one of spooling and sedimentation.



-31-

46. The method of claim 45 wherein said organic solvent is selected from the group consisting of lower alcohols, acetone, polyethylene glycol and dimethylsulfoxide.



FIG.3




FIG.2D

FIG.2A




FIG.1D




FIG.1A



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/10, C07K 1/14</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 97/05248</b> <b>(43) International Publication Date:</b> 13 February 1997 (13.02.97)
<b>(21) International Application Number:</b> PCT/US96/11875 <b>(22) International Filing Date:</b> 18 July 1996 (18.07.96) <b>(30) Priority Data:</b> 08/509,164                      31 July 1995 (31.07.95)                      US <b>(71)(72) Applicant and Inventor:</b> CHOMCZYNSKI, Piotr [US/US]; 778 Avon Fields Lane, Cincinnati, OH 45229 (US). <b>(74) Agents:</b> AHRENS, Gregory, F. et al.; Wood, Herron & Evans, P.L.L., 2700 Carew Tower, Cincinnati, OH 45202 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 6 March 1997 (06.03.97)
<b>(54) Title:</b> SOLUTION CONTAINING CHAOTROPIC AGENT AND PROCESS USING IT FOR ISOLATION OF DNA, RNA AND PROTEINS  <b>(57) Abstract</b>  Solutions and methods are disclosed for the effective, simple isolation/extraction of DNA, RNA and proteins from a single biological material sample, such as cells, tissues and biological fluids. The preferred solutions include effective amounts of a chaotropic agent(s), buffer, reducing agent, and may or may not include an organic solvent. Genomic DNA and total RNA can be isolated utilizing the solutions and methods of the invention in as little as 20 minutes, and proteins in as little as 30 minutes.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/11875

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/10 C07K1/14

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,5 128 247 (KOLLER CHARLES A) 7 July 1992 see column 9 - column 10 ---	1-27, 40-46
X	ANALYTICAL BIOCHEMISTRY, vol. 207, no. 1, 15 November 1992, US, pages 197-201, XP000611804 NELSON J.E. AND KRAWETZ S.A.: "Purification of cloned and genomic DNA by guanidine thiocyanate/isobutyl alcohol fractionation" see abstract see page 198, column 1 --- -/--	1-27, 43-46

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*&\* document member of the same patent family

Date of the actual completion of the international search

12 December 1996

Date of mailing of the international search report

29. 01. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Chakravarty, A

# INTERNATIONAL SEARCH REPORT

Int. onal Application No  
PCT/US 96/11875

**C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. MICROBIOL. METHODS, vol. 19, no. 3, 1994, pages 167-172, XP000610898 LEMA M.W ET AL.: "A general method for the extraction of DNA from bacteria" see page 168 ---	1-27,45, 46
X	EP,A,0 544 034 (SIEMENS AG) 2 June 1993 see the whole document ---	1-5,8,9
X	US,A,5 162 507 (WOLFE SIDNEY N ET AL) 10 November 1992 see column 7, line 48 - line 68 ---	1-11
X	WO,A,92 00983 (MICROPROBE CORP) 23 January 1992 see examples 1,4 -----	1-5

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/11875

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5128247	07-07-92	AU-A- 6432490 CA-A- 2064068 EP-A- 0487646 WO-A- 9102740	03-04-91 15-02-91 03-06-92 07-03-91
EP-A-0544034	02-06-93	CA-A- 2083870 JP-A- 5300174	29-05-93 12-11-93
US-A-5162507	10-11-92	AU-B- 622860 AU-A- 1627988 EP-A- 0368857 JP-T- 2503384 NO-B- 176797 WO-A- 8808849 CA-A- 1337671	30-04-92 06-12-88 23-05-90 18-10-90 20-02-95 17-11-88 28-11-95
WO-A-9200983	23-01-92	CA-A- 2087105 EP-A- 0539515 US-A- 5393672 US-A- 5130423	14-01-92 05-05-93 28-02-95 14-07-92